

The X-ray monitor JEM-X

Purpose:

Precise position determination (< 1 arcmin)
Cover low energyspectrum 3 - 30 keV

Properties:

Field-of-view	12 deg diameter
	5 deg FCFOV
Energy resolution	16% @ 7 keV
	10% @ 20 keV

Imaging: Niels J. Westergaard, Niels Lund, Carl Budtz-Jørgensen

Danish Space Research Institute

Spectra: Stefan Larsson

Stockholm University

INTEGRAL with JEM-X




JEMX October 6, 2004

INTEGRAL Science Analysis
Workshop



JEM-X HARDWARE



JEMX October 6, 2004

INTEGRAL Science Analysis
Workshop

Modes of operation

Full imaging

Restricted imaging

FULL

REST

Rudimentary energy binning, limited time res.

Non-imaging formats (all detector):

Spectral format

SPEC

Timing format

TIME

Spectral timing

SPTI

Grey filter event rejection

When the event buffer is filled above certain level a random rejection of events will be initiated autonomously

Level G will reject a fraction $(31 - G)/32$

$$0 \leq G \leq 31$$

When buffer filling is low again the grey filter level will increase again.

Image production

Steps in the process:

Level	Function	Executable
BIN_1	Shadowgram building	j_ima_shadowgram
BKG	Background subtraction	(included)
IMA	Image reconstruction	j_ima_basic_recon
IMA	Flatfielding	(included)
IMA	Source finding	j_ima_src_find
IMA	Vignetting correction	j_ima_cor_intensity
IMA2	Mosaicking	j_ima_mosaic (OSA5)

Shadowgram building

Event selection based on **STATUS** flag (rowSelect)

Correction for deadtime


Correction for grey filter

$TELAPSE = T_{start} - T_{stop}$

$ONTIME = GTI_1 + GTI_2 + \dots$



$EXPOSURE = ONTIME * corfactor$

shdType = 0 'skew' shadowgram
shdType = 1 standard shadowgram





Mask pattern

25% HURA - but almost not cyclic, hence ghosts almost eliminated




The soon to disappear skew shadowgram




Mask and shadowgram

Mask array from IMOD GRP



Shadowgram from OSA4
shdType=0



Algorithm

Balanced correlation

For each relative position of detector and mask:

Count open and closed mask cells 'on the detector' (N_{open} and N_{closed})

Sum number of counts corresponding to open and closed cells


$$t = \frac{N_{\text{open}}}{N_{\text{open}} + N_{\text{closed}}} \quad (1)$$

Balanced reconstruction:

$$I = (1 - t) \sum S_{\text{open}} - t \sum S_{\text{closed}} \quad (2)$$

Leaking calibration sources



Shadowgram
Crab Nebula
on axis
Rev 102
3 - 6 keV



Shadowgram low, medium energy

3-6 keV, leak removed.



6-12 keV, leak removed




Shadowgram, high energy

12-25 keV

Smoothed version



Background spectrum





Background (1)

3 - 7 keV

30,000 s

7 - 13 keV





Background (2)

13 - 40 keV


30,000 s

radiusLimit = 105




Basic reconstruction example

OSA4 result
Rev 53, PID 66
6-10 keV
Galactic Center region



Fine resolution example

Rev 53
PID 66
Galactic center
6-10 keV



Vignetting function


Vignetting array from
IMOD-GRP
This is the average
vignetting

BUT

complicated geometry
between mask and
collimator and detector

More work!!

Contours: 20, 40, 60, and 80%



New member of ISSW family

Midisky - to become **j_ima_iros** with OSA5



- Backprojection for source finding
- Iterative removal of sources
- As complete modelling of detector as possible including collimator imperfections
- The spectrum and lightcurve extraction based on this modelling will be included in the ISSW as well

Now: `ftp.dsri.dk/pub/jemx/midisky/midi/midi_distribution_2.tar.gz`

Midisky example GRB040812


Full science window

20 s during burst



Midisky manipulated mosaic

In IROS each source gets a position and a strength. Here they are taken out, the remainder smoothed; then the sources have been put back in.



Fin

After running `jemx_science_analysis` up to and including level **IMA** you'll find in the `.../res` directory:


`jmx1_evts_shd.fits`
`jmx1_rsti_ima.fits`
`jmx1_sky_ima.fits`
`jmx1_find_res.fits`
`jmx1_srcl_res.fits`

shadowgrams
'skew' images
WCS images
individual find results
source list summary

and the stage is set for the spectral/timing analysis ...

Crab on axis in log scale

Rev 102
Log scale
3-6 keV



Noter: hvad skal findes

Billeder at skaffe:

Synsfelter

Standardbillede

Fine resolution

midisky = j_ima_iros